

<https://helda.helsinki.fi>

Cadophora margaritata sp. nov. and other fungi associated with the longhorn beetles Anoplophora glabripennis and Saperda carcharias in Finland

Linnakoski, Riikka

2018-11

Linnakoski , R , Kasanen , R , Lasarov , I , Marttinen , T , Oghenekaro , A , Sun , H , Asiegbu , F O , Wingfield , M J , Hantula , J & Heliövaara , K T 2018 , ' Cadophora margaritata sp. nov. and other fungi associated with the longhorn beetles Anoplophora glabripennis and Saperda carcharias in Finland ' , Antonie van Leeuwenhoek , vol. 111 , no. 11 , pp. 2195-2211 . <https://doi.org/10.1007/s10482-018-1112-y>

<http://hdl.handle.net/10138/303100>

<https://doi.org/10.1007/s10482-018-1112-y>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

ArticleTitle	<i>Cadophora margaritata</i> sp. nov., and other fungi associated with the longhorn beetles <i>Anoplophora glabripennis</i> and <i>Saperda carcharias</i> in Finland	
Article Sub-Title		
Article CopyRight	Springer International Publishing AG, part of Springer Nature (This will be the copyright line in the final PDF)	
Journal Name	Antonie van Leeuwenhoek	
Corresponding Author	Family Name	Linnakoski
	Particle	
	Given Name	Riikka
	Suffix	
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	Helsinki, Finland
	Division	
	Organization	Natural Resources Institute Finland (Luke)
	Address	Helsinki, Finland
	Division	Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI)
	Organization	University of Pretoria
	Address	Pretoria, South Africa
	Phone	+358 29 532 2289
	Fax	
Author	Email	riikka.linnakoski@luke.fi
	URL	
	ORCID	http://orcid.org/0000-0002-3294-8088
	Family Name	Kasanen
	Particle	
	Given Name	Risto
	Suffix	
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	Helsinki, Finland
	Phone	
	Fax	
	Email	
	URL	
	ORCID	http://orcid.org/0000-0002-9948-2200
Author	Family Name	Lasarov
	Particle	
	Given Name	Ilmeini

Suffix
Division School of Forest Sciences
Organization University of Eastern Finland
Address Joensuu, Finland
Phone
Fax
Email
URL
ORCID

Author	Family Name	Marttinen
	Particle	
	Given Name	Tiia
	Suffix	
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	Helsinki, Finland
	Phone	
	Fax	
	Email	
	URL	
	ORCID	

Author	Family Name	Oghenekaro
	Particle	
	Given Name	Abbot O.
	Suffix	
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	Helsinki, Finland
	Division	Department of Plant Biology and Biotechnology
	Organization	University of Benin
	Address	Benin City, Nigeria
	Phone	
	Fax	
	Email	
	URL	
	ORCID	

Author	Family Name	Sun
	Particle	
	Given Name	Hui
	Suffix	
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	Helsinki, Finland
	Phone	
	Fax	

Email
URL
ORCID

Author	Family Name	Asiegbu
	Particle	
	Given Name	Fred O.
	Suffix	
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	Helsinki, Finland
	Phone	
	Fax	
	Email	
	URL	
	ORCID	http://orcid.org/0000-0003-0223-7194

Author	Family Name	Wingfield
	Particle	
	Given Name	Michael J.
	Suffix	
	Division	Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI)
	Organization	University of Pretoria
	Address	Pretoria, South Africa
	Phone	
	Fax	
	Email	
	URL	
	ORCID	http://orcid.org/0000-0001-9346-2009

Author	Family Name	Hantula
	Particle	
	Given Name	Jarkko
	Suffix	
	Division	
	Organization	Natural Resources Institute Finland (Luke)
	Address	Helsinki, Finland
	Phone	
	Fax	
	Email	
	URL	
	ORCID	http://orcid.org/0000-0002-1016-0636

Author	Family Name	Heliövaara
	Particle	
	Given Name	Kari
	Suffix	
	Division	Department of Forest Sciences

Organization	University of Helsinki
Address	Helsinki, Finland
Phone	
Fax	
Email	
URL	
ORCID	http://orcid.org/0000-0003-0657-8885

Schedule	Received	25 January 2018
	Revised	
	Accepted	7 June 2018

Abstract Symbiosis with microbes is crucial for survival and development of wood-inhabiting longhorn beetles (Coleoptera: Cerambycidae). Thus, knowledge of the endemic fungal associates of insects would facilitate risk assessment in cases where a new invasive pest occupies the same ecological niche. However, the diversity of fungi associated with insects remains poorly understood. The aim of this study was to investigate fungi associated with the native large poplar longhorn beetle (*Saperda carcharias*) and the recently introduced Asian longhorn beetle (*Anoplophora glabripennis*) infesting hardwood trees in Finland. We studied the cultivable fungal associates obtained from *Populus tremula* colonised by *S. carcharias*, and *Betula pendula* and *Salix caprea* infested by *A. glabripennis*, and compared these to the samples collected from intact wood material. This study detected a number of plant pathogenic and saprotrophic fungi, and species with known potential for enzymatic degradation of wood components. Phylogenetic analyses of the most commonly encountered fungi isolated from the longhorn beetles revealed an association with fungi residing in the *Cadophora*–*Mollisia* species complex. A commonly encountered fungus was *Cadophora spadicea*, a recently described fungus associated with wood-decay. In addition, a novel species of *Cadophora*, for which the name *Cadophora margaritata* sp. nov. is provided, was isolated from the colonised wood.

Keywords (separated by '-') 1 New taxon - Alien invasive species - *Cadophora* sp. - Introduced species - Insect–fungus symbiosis - Longhorn beetles - Vectored pathogen

Footnote Information **Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s10482-018-1112-y>) contains supplementary material, which is available to authorized users.

2 ***Cadophora margaritata* sp. nov., and other fungi associated**
3 **with the longhorn beetles *Anoplophora glabripennis***
4 **and *Saperda carcharias* in Finland**

5 Riikka Linnakoski  · Risto Kasanen  · Ilmeini Lasarov · Tiia Marttinen ·
6 Abbot O. Oghenekaro · Hui Sun · Fred O. Asiegbu  · Michael J. Wingfield  ·
7 Jarkko Hantula  · Kari Heliövaara 

8 Received: 25 January 2018 / Accepted: 7 June 2018
9 © Springer International Publishing AG, part of Springer Nature 2018

10 **Abstract** Symbiosis with microbes is crucial for
11 survival and development of wood-inhabiting long-
12 horn beetles (Coleoptera: Cerambycidae). Thus,
13 knowledge of the endemic fungal associates of insects
14 would facilitate risk assessment in cases where a new
15 invasive pest occupies the same ecological niche.
16 However, the diversity of fungi associated with insects
17 remains poorly understood. The aim of this study was
18 to investigate fungi associated with the native large
19 poplar longhorn beetle (*Saperda carcharias*) and the
20 recently introduced Asian longhorn beetle (*Ano-*
21 *plophora glabripennis*) infesting hardwood trees in
22 Finland. We studied the cultivable fungal associates

obtained from *Populus tremula* colonised by *S.* 23
carcharias, and *Betula pendula* and *Salix caprea* 24
infested by *A. glabripennis*, and compared these to the 25
samples collected from intact wood material. This 26
study detected a number of plant pathogenic and 27
saprotrophic fungi, and species with known potential 28
for enzymatic degradation of wood components. 29
Phylogenetic analyses of the most commonly encoun- 30
tered fungi isolated from the longhorn beetles revealed 31
an association with fungi residing in the *Cadophora*– 32
Mollisia species complex. A commonly encountered 33
fungus was *Cadophora spadicis*, a recently described 34
fungus associated with wood-decay. In addition, a 35
novel species of *Cadophora*, for which the name 36
Cadophora margaritata sp. nov. is provided, was 37
isolated from the colonised wood. 38

A1 **Electronic supplementary material** The online version of
A2 this article (<https://doi.org/10.1007/s10482-018-1112-y>) con-
A3 tains supplementary material, which is available to authorized
users.

A4 R. Linnakoski (✉) · R. Kasanen · T. Marttinen ·
A5 A. O. Oghenekaro · H. Sun · F. O. Asiegbu ·
A6 K. Heliövaara
A7 Department of Forest Sciences, University of Helsinki,
A8 Helsinki, Finland
A9 e-mail: riikka.linnakoski@luke.fi

A16 I. Lasarov
A17 School of Forest Sciences, University of Eastern Finland,
A18 Joensuu, Finland

A19 A. O. Oghenekaro
A20 Department of Plant Biology and Biotechnology,
A21 University of Benin, Benin City, Nigeria

A10 R. Linnakoski · J. Hantula
A11 Natural Resources Institute Finland (Luke), Helsinki,
Finland

A12 R. Linnakoski · M. J. Wingfield
A13 Department of Microbiology and Plant Pathology,
A14 Forestry and Agricultural Biotechnology Institute (FABI),
A15 University of Pretoria, Pretoria, South Africa

Keywords 1 New taxon · Alien invasive species ·
Cadophora sp. · Introduced species · Insect–fungus
 symbiosis · Longhorn beetles · Vectored pathogen

Introduction

A nutrient-poor wood tissue (xylem), lacking readily available carbohydrates and proteins, is a challenging niche to be exploited by other organisms. However, cerambycid beetles (Coleoptera: Cerambycidae) are adapted to complete their long lifecycle in wood tissues. In this niche, symbiosis with microbes (including filamentous fungi and yeasts) has been considered to be crucial for their survival and development (Kukor et al. 1988; Schloss et al. 2006; Watanabe and Tokuda 2010; Scully et al. 2013). The role of the associated microbes can be considered as directly nutritional, in which insect larvae cultivate and feed on the fungi (Francke-Grosmann 1967), or when either ingested fungal enzymes (Kukor and Martin 1986; Kukor et al. 1988) or endosymbiotic gut microbes contribute to cellulose degradation (Uvarov 1929; Kukor et al. 1988; Grünwald et al. 2010; Watanabe and Tokuda 2010; Scully et al. 2013). These symbiotic fungi may also have other functional roles, such as nitrogen and vitamin acquisition (Gibson and Hunter 2009; Ayayee et al. 2014).

The association of Cerambycidae and fungi was recognised relatively long ago (Uvarov 1929; Jurzitza 1959; Riba 1977; Chararas and Pignal 1981; Nardon and Grenier 1989). In general, the modified colour and texture of the wood close to the larval tunnels indicate that symbiotic organisms are commonly involved in the diet of xylophagous insects. Many of the wood-inhabiting fungi and insect symbioses have long co-evolutionary histories (Vega and Blackwell 2005), recently supported by the observations of fungi and associated organisms successfully passing through the insect alimentary tract (Drenkhan et al. 2013, 2016). The associations with Cerambycidae have arisen independently in distantly related fungal genera, representing examples of convergent evolution (Jones et al. 1999). Cerambycidae are known to harbour a rich assemblage of microbes in terms of taxonomic and functional diversity (Grünwald et al. 2010; Watanabe and Tokuda 2010; Scully et al. 2013; Ayayee et al.

2014). However, the diversity of fungi associated with most Cerambycidae remains to be investigated.

An example of a common but poorly known Cerambycidae is the large poplar longhorn (*Saperda carcharias*) (Fig. 1). The biology of the insect is not fully understood, and there are no previous reports of its symbiotic fungi. The beetle is considered as the most harmful pest of hybrid poplars and aspen (*Populus tremula*) in Finland (Hallaksela 1999; Välimäki and Heliövaara 2007). The adult beetle feeds on aspen leaves and lays eggs in basal parts of the stems during August–September (Heliövaara et al. 2004). The eggs undergo overwintering for 10–11 months and the emerging larvae remain in the tunnels for 2–4 years. During this period, larvae can gnaw tunnels up to 1 m long in the xylem. These tunnels are often surrounded by a characteristic dark brown colour expanding vertically 1–3 m, which prevent the logs from being used in the mechanical wood industry. Part of the wood material gnawed by the larvae is transported through their alimentary canals. Due to long larval period of development in the tunnels, it is likely that the poplar longhorn has interactions with a number of fungi during its life cycle.

The Asian longhorn beetle (*Anoplophora glabripennis*) is a serious forest pest native to China (Lingafelter and Hoebeke 2002) (Fig. 2). In its native range, the beetle is not considered as a pest in natural forests. However, the beetle became a common pest in China as a result of widespread planting of susceptible poplar (*Populus* sp.) hybrids (EPPO 2016). As an invasive species, *A. glabripennis* is a highly destructive quarantine pest (EPPO 2016). The beetle is capable of infesting a wide range of deciduous trees, including also healthy trees. The Asian longhorned beetle has spread globally mainly in ineffectively treated wooden packaging materials originating from China. It was accidentally introduced first into North America (Haack et al. 1996; Hu et al. 2009), followed by several introductions into European countries during the past two decades (Tomiczek et al. 2002; Hérard et al. 2006, 2009; Haack et al. 2010), and recently into Finland.

In several instances, baseline information on fungal associates of insects in their endemic environment is lacking. This hinders risk assessment measures in cases where a novel forest pest and/or pathogen is introduced. Novel insect–fungal interactions represent



Fig. 1 **a** Aspen (*P. tremula*) stand in Urjala, Finland, **b**, **c** the large poplar longhorn beetle (*S. carcharias*) adult and larvae, and **d** brown decay surrounding the beetle gallery on *P. tremula*

an increased risk to forest health (Humble and Allen 2006; Hulcr and Dunn 2011; Wingfield et al. 2016; Stenlid and Oliva 2016). Examples of devastating tree disease problems that have arisen from these associations are increasing in number (Lu et al. 2011; Akbulut and Stamps 2012; Ploetz et al. 2013; Santini and Faccoli 2014; Wingfield et al. 2016). Recent introductions of plant pathogens (Wingfield et al. 2016) suggest that latent microbial pathogens commonly remain undetected in quarantine inspections.

The aim of the study was to provide baseline information on fungal diversity associated with *S.*

carcharias in its native host tree and environment, and with the introduced *A. glabripennis* infestation in Finland. Our hypotheses were that the longhorn beetles are associated with specific fungi, which may improve nutritional quality of the larvae facilitating larval development inside the wood. Furthermore, we hypothesized that differences in fungal species compositions between intact wood and wood colonized by beetles would reflect the specific symbiotic fungi associated with each beetle. In addition, we hypothesize that the introduced insect pest would establish interactions with native fungal associates of



Fig. 2 The Asian longhorned beetle (*A. glabripennis*) adult and larvae, and brown decay surrounding the beetle gallery on *B. pendula*

ecologically and taxonomically closely related insect species.

Materials and methods

Study areas and collection of samples

Populus tremula trees damaged by *S. carcharias* were collected from a 15-year-old aspen stand in Urjala, southwestern Finland (66°00'N, 23°29'E) in May 2013 (Fig. 1). The aspen clonal stand 2120/01 was established on 25 May 1998 as a part of the aspen clonal research programme of the Finnish Forest Research Institute. The average diameter at breast height of trees was 12 cm and their average height was ca. 9.0 m. The stand consisted of both aspen and hybrid aspen, which were planted at intervals of 3 m in 3.0 × 3.0 m blocks. All the five sampled trees were of the same clone, collected from different blocks. The logs including the entire beetle tunnels were cut with a chain saw immediately closed in plastic bags and stored at + 4 °C.

The logs (60–90 cm of length) were sawn with table mounted circular saw longitudinally into two halves in a laboratory for fungal isolations. Two samples 5 mm × 10 mm (separated by 5 cm) from three points (separated by at least 15 cm) were cut from the brown-coloured wood adjacent to larval

tunnels with surface disinfected carpenter's knife and surgical knife. Control isolations were made from the non-coloured, undamaged distal ends of the logs as described above. All samples were surface disinfected in sterile conditions with 70% ethanol (10 s) and 10% sodium hypochlorite (NaClO) (5 s) and finally washed four times with fresh sterile water. The samples were placed onto both water agar (WA: 2% AMRESCO® agar, bacteriological grade) and malt extract agar (MEA: 2% Bacto™ malt extract agar and 1% AMRESCO® agar, bacteriological grade) in 90 mm Petri dishes. A subset of unsterilised wood samples was incubated at 25 °C on sterilised filter paper placed in clean Petri dishes (9 cm).

Anoplophora glabripennis-infested trees were detected in Vantaa, Finland (60.303'N, 25.119'E) in October–November 2015. During the inspections and eradication measures, a total of 20 *Betula pendula* and 13 *Salix caprea* trees showing signs of *A. glabripennis* infestation were detected (Fig. 2). The trees were removed by the Finnish Food Safety Authority (Evira), and transported to a quarantine facility for careful examination and destruction. At the same time, fungal isolations were made from decayed wood surrounding the galleries of *A. glabripennis* on five *B. pendula* and two *S. caprea* trees, using a similar sampling scheme as described earlier. Isolations were also made from the surfaces of five living larvae found in galleries on *S. caprea*. Small pieces of wood and the

larvae were placed onto the surface of WA and MEA. The larvae were removed from the plates after approximately 15 min. Control isolations were made from the non-coloured, undamaged distal ends of the logs.

The cultures obtained from *S. carcharias* and *A. glabripennis* were grown for 1 week in the dark at + 25 °C and inspected daily. Emerging mycelia was sub-cultured to obtain pure cultures. Purified fungal isolates were grown on MEA and grouped based on their colony and morphological characteristics. Representative isolates from each group were chosen for identification based on DNA sequences. Isolates were deposited in the Culture Collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa, and the ex-type isolates also in the CBS-KNAW Collections, Westerdijk Fungal Biodiversity Institute, Utrecht, the Netherlands (Table 1). Herbarium specimens were deposited in the Herbarium of the University of Turku, Finland (TUR), Finland.

DNA extraction, PCR and sequencing

Pure cultures for DNA extractions were cultivated on MEA plates. DNA was extracted from at least one isolate of each morphological group using the methods described previously (Terhonen et al. 2011; Linnakoski et al. 2016). Gene regions sequenced in the study were the internal transcribed spacer (ITS) regions (ITS1 and ITS2) including the 5.8S gene (all samples), as well as the partial beta-tubulin gene (only for the novel species).

The DNA samples from mycelial cultures obtained from *S. carcharias* were amplified using a PCR protocol for DyNAzyme DNA[®] polymerase (Finnzymes, Espoo, Finland). A PCR reaction (total volume 50 µl) contained 30.5 µl of sterile water, 5 µl of DyNAzyme 10 × buffer, 1 µl of 10 mM dNTP's, 1 µl of 25 mM ITS1-F primer (Gardes and Bruns 1993), 1 µl of 25 mM ITS4 primer (White et al. 1990), 1 µl of 50 mM MgCl₂, 0.5 µl of DyNAzyme DNA[®] polymerase and 10 µl/l of DNA template. The partial beta-tubulin gene region was amplified using the primer pair T10 (O'Donnel and Cigelnik 1997) and Bt2b (Glass and Donaldson 1995). The PCR reactions were run with BIOER XP Cycler (BIOER Technology, Hangzhou, China). The PCR reaction was started with 4 min of initial denaturation followed by 30 cycles of

45 s of denaturation (95 °C), 45 s of annealing 52 °C, and 1 min extension (72 °C). A final extension step was 10 min (72 °C). Products were visualised by electrophoresis on a 1.5% agarose gel stained with ethidium bromide and purified using GenElute[™] PCR clean-up kit according to the manufacturer's instructions (Sigma-Aldrich, Missouri, USA).

Amplification of the ITS region and purification of the PCR products of fungal samples from *A. glabripennis* were performed following the same protocols described previously (Linnakoski et al. 2016). The studied gene region was amplified using the primer pair ITS1-F (Gardes and Bruns 1993) and ITS4 (White et al. 1990). The reaction mixture contained 0.15 µl of MyTaq[™] DNA Polymerase (5 U/µl) (Bioline, Massachusetts, USA), 5 µl of MyTaq[™] Reaction Buffer (5 ×) containing dNTPs, MgCl₂ and enhancers, 0.50 µl of each primer (10 mM stock concentration) (Whitehead Scientific Ltd, Cape Town, South Africa), and 1 µl fungal genomic DNA. Sterile Sabax water was added to adjust the reaction volume to 25 µl. PCR amplifications were performed using the following conditions: denaturation at 95 °C for 2 min, followed by 32 cycles of 30 s at 94 °C, 30 s at 55 °C and 1 min at 72 °C, and a final extension at 72 °C for 7 min. An aliquot of 5 µl of the each PCR product was stained with GelRed[™] nucleic acid gel stain (Biotium, Hayward, CA, USA), run on 2% agarose gel at 60 V along with a 100 bp molecular marker (Fermentas O' Gene Ruler[™]) and visualised with a Gel Doc EZ Imager (Bio-Rad Laboratories Inc.). Amplified PCR products were cleaned using the Exonuclease I—Shrimp Alkaline Phosphatase Clean Up (Exo-SAP) protocol.

The sequencing reactions contained 0.5 µl of BigDye[®] Terminator v3.1 Ready Reaction mixture (Perkin-Elmer Applied Biosystems, Warrington, UK), 2.1 µl of sequencing buffer, 1 µl of either the forward or reverse primer (10 mM stock concentration) and 1 µl of purified PCR product. Sterile Sabax water was added to adjust the reaction volume to 12 µl. The thermal cycling conditions were: 25 cycles of 10 s at 96 °C, 5 s at 55 °C and 4 min at 60 °C. Sequencing products were then cleaned using ethanol/salt precipitation. Sequencing of the isolates from *S. carcharias* was conducted on the Applied Biosystems (ABI) with an ABI 3130xl sequencer (Thermo Fisher Scientific, Casrbad, USA) at the Haartman Institute Sequencing Unit, University of Helsinki. Sequencing of the



Table 1 Fungal isolates obtained from longhorn beetles *A. glabripennis* and *S. carcharias* colonized wood and intact wood in Finland

Species	Culture collection no. ^a	Herbarium no. ^b	Beetle	Host tree	Substrate	Date of isolation	GenBank acc. no.	
							LSU	ITS
<i>Alternaria</i> sp.	N/A		<i>Saperda carcharias</i>	<i>Populus tremula</i>	Colonized wood	01 Aug 2013		KJ702019
<i>Arthrinium</i> sp.	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	20 Sept 2013		KJ702039
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	20 Sept 2013		KJ702044
<i>Ascochyne cylindrium</i>	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Aug 2013		KJ702030
<i>Cadophora spadicis</i>	CMW49956		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015		MF188968
	CMW49957		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	09 Dec 2015		MF188969
	CMW49958		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	09 Dec 2015		MF188970
	CMW49959		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	09 Dec 2015		MF188971
	CMW49960		<i>A. glabripennis</i>	<i>Salix caprea</i>	Larvae	11 Dec 2015		MF188972
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Sept 2013		KJ702035
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Sept 2013		KJ702037
<i>Cadophora margaritata</i> sp. nov.	CMW51780, CBS144083 ^c	TUR207199 ^e	<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013	MH267288	KJ702027
	CMW51781, CBS144084 ^d	TUR207200 ^d	<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013		MH203866
<i>Cladosporium</i> sp.	CMW49961		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015		MF188973
	CMW49962		<i>A. glabripennis</i>	<i>S. caprea</i>	Larvae	11 Dec 2015		MF188974
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013		KJ702022
<i>Coniochaeta</i> sp.	N/A		<i>Anoplophora glabripennis</i>	<i>Betula pendula</i>	Colonized wood	23 Nov 2015		MH203865
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Aug 2013		KJ702032
<i>Coniothyrium carteri</i>	49963		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015		MF188975
<i>Cosmospora</i> sp.	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	02 Aug 2013		KJ702026

Table 1 continued

Species	Culture collection no. ^a	Herbarium no. ^b	Beetle	Host tree	Substrate	Date of isolation	GenBank acc. no.	
							LSU	ITS
<i>Epicoecum</i> sp.	CMW49964		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015	MF188976	MH327788
	CMW49965		<i>A. glabripennis</i>	<i>S. caprea</i>	Larvae	11 Dec 2015	MF188977	MH327789
<i>Fusarium</i> sp.	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Intact wood	18 Nov 2013	KJ702040	
	CMW49966		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015	MF188978	
	CMW49967		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015	MF188979	MH327790
	CMW49968		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015	MF188980	
	CMW49969		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015	MF188981	
	CMW49970		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	09 Dec 2015	MF188982	MH327791
	CMW49971		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	09 Dec 2015	MF188983	
	CMW49972		<i>A. glabripennis</i>	<i>S. caprea</i>	Colonized wood	11 Dec 2015	MF188984	MH327792
	N/A		<i>A. glabripennis</i>	<i>S. caprea</i>	Intact wood	11 Dec 2015	MH203867	
	N/A		<i>A. glabripennis</i>	<i>S. caprea</i>	Colonized wood	11 Dec 2015	MF188985	
<i>Mollisia dextrinospora</i>	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	02 Aug 2013	KJ702020	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Intact wood	18 Nov 2013	KJ702041	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	18 Nov 2013	KJ702045	
	CMW49973		<i>A. glabripennis</i>	<i>B. pendula</i>	Colonized wood	23 Nov 2015	MF188986	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	02 Aug 2013	KJ702018	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Intact wood	18 Nov 2013	KJ702042	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013	KJ702017	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013	KJ702024	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013	KJ702025	
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013	KJ702025	

Table 1 continued

Species	Culture collection no. ^a	Herbarium no. ^b	Beetle	Host tree	Substrate	Date of isolation	GenBank acc. no.	
							LSU	ITS β -Tubulin
<i>Phialocephala lagerbergii</i>	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Sept 2013		KJ702036
<i>Phlebia radiata</i>	CMW49974		<i>A. glabripennis</i>	<i>S. caprea</i>	Colonized wood	11 Dec 2015		MF188987
<i>Pholiota</i> sp.	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013		KJ702028
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Aug 2013		KJ702029
<i>Phoma</i> sp.	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	20 Sept 2013		KJ702031
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	20 Sept 2013		KJ702034
<i>Physalospora scirpi</i>	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	13 Aug 2013		KJ702033
<i>Pseudeurotium bakeri</i>	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013		KJ702021
	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	01 Aug 2013		KJ702023
<i>Sarocladium strictum</i>	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Intact wood	18 Nov 2013		KJ702043
<i>Tolypocladium</i> sp.	N/A		<i>A. glabripennis</i>	<i>S. caprea</i>	Colonized wood	11 Dec 2015		MH203868
<i>Trichoderma</i> sp.	N/A		<i>S. carcharias</i>	<i>P. tremula</i>	Colonized wood	20 Sept 2013		KJ702038

Bold *Cadophora-Mollisia* species complex

^aCMW, Culture Collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa; CBS, the CBS-KNAW Collections, Westerdijk Fungal Biodiversity Institute, Utrecht, the Netherlands

^bTUR Herbarium, Centre for Biodiversity, University of Turku, Finland

^cEx-type

^dEx-paratype

^eHolotype

isolates from *A. glabripennis* was conducted on the ABI with an ABI 3500xl sequencer (Thermo Fisher Scientific, Carlsbad, USA) at the DNA Sequencing Facility of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria.

Sequence analyses and fungal identification

Consensus sequences were assembled with the Geneious R6.0.6 (Biomatters Ltd, Auckland, New Zealand), after which initial identification of the isolates was obtained using a BLAST search in GenBank nucleotide database (<http://www.ncbi.nlm.nih.gov>) applying a megablast algorithm and (a) excluding uncultured/environmental sample sequences and (b) limiting the search on sequences from type material (Supplementary Table 1). Identification to the closest genus and species level was made as far as possible, with caution given to the fact that the ITS sequence variability in certain fungal groups is low, and acknowledging possibly misidentified sequences in GenBank. We considered reliable identification to consist of the BLAST matches that had $\geq 98\%$ sequence similarity to ex-type sequences or peer-reviewed published studies. For species in the *Cadophora-Mollisia* complex, BLAST searches were also used to retrieve similar sequences, as well as from type material to compile data sets for phylogenetic analysis. Data sets were edited with MEGA v.7 (Kumar et al. 2016). GenBank accession numbers of sequences are presented in the corresponding phylogenetic trees (Figs. 3, 4).

Cadophora finlandica and *Melinomyces variabilis* were used as outgroups for the ITS data set, and *Cadophora luteo-olivacea* and *Cadophora malorum* for the beta-tubulin. The data sets were aligned using the online version of MAFFT v. 7 (Kato and Standley 2013). Three different phylogenetic methods were applied: maximum likelihood (ML), maximum parsimony (MP), and Bayesian inference (BI). ML was performed in the online version of PhyML 3.0 (Guindon et al. 2010), using automatic model selection by SMS (Lefort et al. 2017) and Akaike information criterion (AIC) (Sugiura 1978). Confidence levels were estimated with 1000 bootstrap replicates. MP analyses were conducted using PAUP v. 4.0b10 (Swofford 2002). Gaps and missing data were excluded in the MP analyses. For BI analyses, the best-fitting evolutionary model using MrModeltest 2.3

(Nylander 2004) based on the Akaike Information Criterion (AIC) were determined. BI analyses based on a Markov Chain Monte Carlo (MCMC) simulation were carried out with MrBayes v3.2.2 (Ronquist and Huelsenbeck 2003). The MCMC chains were run for five million generations using a sample frequency of 100 (resulting in 50,000 trees). Burn-in values were determined for the respective data sets, and all sampled trees having lower than the burn-in values were discarded. The resulting majority rule consensus trees were viewed with MEGA v.7 (Kumar et al. 2016), and post-processed with Adobe Illustrator CC 2018 (Adobe, San Jose, USA).

Results

Isolation and identification of fungi

Fungi were isolated from all of the collected samples associated with *S. carcharias* and *A. glabripennis* infestations in Finland. The study resulted in total of 114 fungal isolates. The trimmed consensus sequences for the ITS gene regions ranged between 493 and 574 bp. The sequences obtained in this study were deposited in GenBank and their accession numbers are presented in the Table 1.

The fungal species richness was higher in the colonised parts than in the intact parts of the wood (Table 2). The most common fungi isolated from both *A. glabripennis* and *S. carcharias*-colonised wood were members in the *Cadophora-Mollisia* species complex. These fungi were not found in the intact parts of the wood. Only a small number of fungi were detected in both the colonised and intact wood samples, including *Epicoccum* sp. and *Fusarium* sp. *Fusarium* sp. was the most commonly isolated species in the intact part of *P. tremula* sample logs. Some species were detected only from the intact wood including *Mortierella hyalina* and *Sarocladium strictum*.

In total, 83 isolates were obtained from *S. carcharias* in Finland. These represented at least 18 species of Ascomycota and one species of Basidiomycota (Table 1). The most common species isolated from *S. carcharias* colonised wood was *Cadophora spadicea*, a putatively novel *Cadophora* species and *Pseudeurotium bakeri*. The other fungal species were isolated only occasionally.





Fig. 3 Phylogenetic tree of species in the *Cadophora*–*Mollisia* species complex obtained from ML analyses of the ITS data set. Sequences obtained in this study are printed in bold type. Bootstrap support values above 70% for ML/MP are presented

at the nodes. Posterior probabilities (above 70%) obtained from BI are indicated by bold lines at the relevant branching points. *bootstrap values < 70%. T, ex-type isolates. Scale bar = total nucleotide difference between taxa

Fig. 4 Phylogenetic tree of species in the *Cadophora–Mollisia* species complex obtained from ML analyses of the beta-tubulin data set. Sequences obtained in this study are printed in bold type. Bootstrap support values above 70% for ML/MP are presented at the nodes. Posterior probabilities (above 70%) obtained from BI are indicated by bold lines at the relevant branching points. *bootstrap values < 70%. T, ex-type isolates. Scale bar = total nucleotide difference between taxa



Isolates from *A. glabripennis* in Finland represented at least seven species of Ascomycota and one species of Basidiomycota (Table 1). In addition to the *Cadophora–Mollisia* isolates, other commonly detected fungi from *A. glabripennis*-infested wood and larvae included a *Fusarium* sp. closely related to an extensively studied gut symbiont isolate previously derived from an *A. glabripennis* larvae (Geib et al. 2012; Herr et al. 2016). The other fungi were isolated only occasionally.

DNA sequence analysis

The identities of isolates representing the most commonly encountered genera, *Cadophora–Mollisia*, were further confirmed by phylogenetic analysis. Data originating from the previous studies by Harrington and McNew (2003) and Travadon et al. (2015) served as a backbone for the phylogenetic tree construction for the *Cadophora* and allied species.

The aligned ITS data set included 44 taxa and 575 characters (with gaps). The best-fitting substitution model estimated for the *Cadophora* ITS data set was GTR + I + G. The ML, MP and BI analyses produced trees with similar topologies (Fig. 3).

The ITS data were sufficient to distinguish between most species included in the data set. However, it did not distinguish between *Cadophora melinii* and *Cadophora fastigiata*, and species in the *Phialocephala lagerbergii* clade. Phylogenetic analyses for *Cadophora–Mollisia* confirmed the identities of four species isolated in this study. These were *C. spadicis*, *P. lagerbergii*, *Mollisia dextrinospora*, and a putatively novel species (Fig. 3). The identity of the unknown species was further confirmed by phylogenetic analysis of sequences for the partial beta-tubulin gene region.

The aligned beta-tubulin data set for the *Cadophora* included 18 taxa and 523 characters (with gaps). The best-fitting substitution model estimated for the data set was GTR + I + G. The novel species was clearly distinct from the other currently known species of *Cadophora* (Fig. 4). The phylogenetic analysis also supported the recent phylogenetic placement of *Cadophora* in four separate clades (Travadon et al. 2015).

Taxonomy

Based on the phylogenetic analyses for ITS region and partial beta-tubulin gene, as well as the morphological

Table 2 Number of fungal isolates obtained from *A. glabripennis* and *S. carcharias* in Finland

Species	<i>Anoplophora glabripennis</i>		<i>Saperda carcharias</i>		Total
	Colonized wood	Intact wood	Colonized wood	Intact wood	
<i>Alternaria</i> sp.	0	0	4	0	4
<i>Arthrimum</i> sp.	0	0	2	2	4
<i>Ascocoryne cylichnium</i>	0	0	2	0	2
<i>Cadophora spadicis</i>	7	0	7	0	14
<i>Cadophora margaritata</i> sp. nov.	0	0	10	0	10
<i>Cladosporium</i> sp.	3	0	4	0	7
<i>Coniochaeta</i> sp.	1	0	7	0	8
<i>Coniothyrium carteri</i>	1	0	0	0	1
<i>Cosmospora</i> sp.	0	0	7	0	7
<i>Epicoccum</i> sp.	2	0	1	1	4
<i>Fusarium</i> sp.	12	1	1	4	18
<i>Mollisia dextrinospora</i>	1	0	0	0	1
<i>Mortierella gamsii</i>	0	0	1	0	1
<i>Mortierella</i> cf. <i>hyalina</i>	0	0	0	2	2
<i>Mucor</i> sp.	0	0	6	0	6
<i>Phialocephala lagerbergii</i>	0	0	2	0	2
<i>Phlebia radiata</i>	1	0	0	0	1
<i>Pholiota</i> sp.	0	0	5	0	5
<i>Phoma</i> sp.	0	0	2	0	2
<i>Physalospora scirpi</i>	0	0	1	0	1
<i>Pseudeurotium bakeri</i>	0	0	10	0	10
<i>Sarocladium strictum</i>	0	0	0	2	2
<i>Tolypocladium</i> sp.	0	1	0	0	1
<i>Trichoderma</i> sp.	0	0	1	0	1
Total	28	2	73	11	114

characteristics and ecology, the isolates of one of the *Cadophora* sp. isolated formed a distinct lineage that represents a novel taxon. This is described as follows with measurements presented as (min–)(mean – SD)–(mean + SD)(–max)’.

***Cadophora margaritata* R. Linnakoski, I. Lasarov & A.O. Oghenekaro sp. nov.** (Fig. 5). MycoBank: MB 825227.

Etymology: The name is derived from the Latin word ‘margaritata’ for ‘beaded’ or ‘pearls’ and refers to beaded appearance of conidial mass.

Sexual state: Not observed.

Asexual state: Aerial mycelium on MEA consisted of pale brown, tuberculate, septate hyphae up to (1.2–)1.5–2.3(–3.2) µm diam. Conidiophores arising from aerial hyphae mostly branched (2–12 branches), pale brown, septate, (93–)114.8–243(–295) µm long

excluding conidia, (1.4–)2.1–3(–3.5) µm wide (Fig. 5a, b). Conidiogenous cells terminal or lateral, mono- or poly-phialidic, hyaline (or paler brown), (8.9–)12.3–22(–29.2) µm long, slightly swollen at base, (1.6–)1.9–2.9(–3.6) µm at the widest part, tapering significantly towards collarette; collarettes cup-shaped, (1.3–)2–3(–3.7) µm wide and (0.8–)1–1.9(–2.4) µm long (Fig. 5b). Conidia aseptate, hyaline, ovoid, truncate at base (1.9–)2.5–3.7(–4.4) long × (1.3–)1.4–2.2(–3.2) µm wide, aggregating into globose (‘beaded appearance’) of conidial mass. **Culture characteristics:** Colonies on 2% MEA olivaceous black (Fig. 5d), mycelium superficial on agar, aerial mycelium sparse. Culture radial growth rate 3.1 mm/d (± 0.3) at 22 °C.

Specimens examined: FINLAND, Western Finland Province: Urjala, isolated from *Saperda carcharias*




Fig. 5 Morphological characteristics of the *C. margaritata* (CMW 51780 = CBS 144083). **a** Conidiophores, **b** conidiogenous cells, **c** conidia, **d** culture on MEA. Scale bars **a–c** 10 µm

infested *Populus tremula*, collector Tiia Martinen, 01 August 2013. Holotype dried specimen TUR 207199 (<http://mus.utu.fi/TFU.207199>), ex-holotype living culture CMW 51780 = CBS 144083; Paratype dried specimen TUR 207200 (<http://mus.utu.fi/TFU.207200>), ex-paratype living culture CMW 51781 = CBS 144084.
Host tree: Populus tremula.
Insect vector: Saperda carcharias.

Discussion

This study contributes to limited investigations on mycobiota associated with cerambycid beetles, and is the first to report fungi associated with the large poplar longhorn (*S. carcharias*). We isolated a number of

plant pathogens, saprobes and species with known potential to cause wood degradation in the galleries of the *S. carcharias* on *P. tremula*. In total 20 fungal species residing in two phyla were identified, the most common detected in the beetle-colonised wood were species of *Cadophora* and allied species. This study also identified fungi associated with material collected from *A. glabripennis* material retained in quarantine in Finland. At least eight fungal species were detected, the most common of these were species of *Cadophora* and *Fusarium*. While *A. glabripennis* is known to harbour endosymbiotic fungi (Scully et al. 2013, 2014), these results are amongst the first to provide evidence of wood-inhabiting fungi associated with *A. glabripennis* in its introduced range. It was also noteworthy that the introduced pest shares a

number of fungal associates with the native species (Table 1).

One of the intriguing findings in this study was the strong presence of a number of *Cadophora* and allied species existing in association with the longhorn beetles. This study is the first to reveal an association of these fungi with insects. Four *Cadophora* and allied species were identified, including the novel species *C. margaritata* described in. *Cadophora* species were originally described as fungi causing blue stain of lumber (Lagerberg et al. 1927). These fungi are now known to reside in two distinct genera *Phialophora* (Chaetothyriales, Eurotiomycetes) and *Cadophora* (Helotiales, Leotiomycetes) (Gams 2000; Harrington and McNew 2003). *Cadophora* has received little attention in the past but recent reports implicating species as potential grapevine pathogens in various countries have changed this view (Halleen et al. 2007; Casieri et al. 2009; Gramaje et al. 2011; Navarrete et al. 2011; Úrbez-Torres et al. 2014). *Cadophora spadici*s detected in the present study is one of the recently described species reported from wood-decay of grapevine in North America (Travadon et al. 2015). A previous study has also reported that *Cadophora* species are able to cause dark staining and extensive soft rot in *Betula* and *Populus* wood in vitro (Blanchette et al. 2004). Based on past findings, it is reasonable to assume that *Cadophora* species reported here play some role in wood degradation close to the larval tunnels of Cerambycidae in general.

The novel species described in this study, is characterised by production of phialides with hyaline collarettes, morphological structures typical for *Cadophora* and allied phialide-producing fungi. An example is *Mollisia dextrinospora*, which has been reported to have a cadophora-like asexual stage (Greenleaf and Korf 1980). As shown in the present study and also in previous phylogenetic analyses (Crous et al. 2003; Harrington and McNew 2003; Day et al. 2012; Pärtel 2016), several species of *Cadophora* and *Mollisia* are closely related. Correct identification of these species is hindered by the occurrence of numerous misidentified entries in nucleotide sequence databases (Pärtel 2016). Consequently, *Mollisia* appears to be a heterogeneous assemblage of fungi that requires taxonomic treatment. The *Mollisia* isolate found in this study grouped with the ex-type strain of *M. dextrinospora* and its identity seems clear. The species was originally described from decayed *Acacia* wood in Madeira

(Greenleaf and Korf 1980), and it could have a similar role in wood decay in its association with the longhorn beetles. The identity of the fourth species belonging in the *Cadophora*–*Mollisia* species complex found in this study, a fungus closely related to *Phialocephala lagerbergii* could not be resolved due to lack of available type sequence data.

Fusarium is a highly diverse cosmopolitan genus that includes plant and human pathogens, saprobes and endophytes. Therefore, it is not surprising that these fungi have also been found as symbionts of cerambycid and ambrosia beetles (Geib et al. 2012; Morales-Ramos et al. 2000; Kasson et al. 2013). The presence of these fungi with *A. glabripennis* in the present study is perhaps not surprising but their consistent occurrence suggests that they may be relevant to *A. glabripennis*. This is supported by the fact that they have previously been found consistently associated with the pest in other areas and in different host trees (Geib et al. 2012). In this association, species of *Fusarium* have been considered as internal gut symbionts supporting cellulose degradation and larval fitness of *A. glabripennis* (Scully et al. 2013, 2014). Species in this group are also symbionts of ambrosia beetles (Morales-Ramos et al. 2000; Kasson et al. 2013). However, in this study *Fusarium* spp. were also detected as the most common isolates in healthy parts of *P. tremula* wood, indicating their role as endophytes on aspen.

An *Aureobasidium* sp. was commonly isolated in this study from *P. tremula*. *Aureobasidium* species appear to be a common associate of aspen *P. tremula* (Santamaría and Diez 2005), with probable fitness effects. Albrechtsen et al. (2010) reported that aspen clonal stands, where *Aureobasidium* sp. was present, were more resistant to browsing by herbivores. Another commonly detected fungus was *Pseudeurotium bakeri* in the *S. carcharias* colonised wood. The fungus has an ability to utilize lignin residues left from brown rot and humus (Ran et al. 2016). *Pseudeurotium bakeri* has also been found as a root-endophyte of the common grass, *Dactylis glomerata* (Sánchez Márquez et al. 2007). The other fungi were detected less commonly. These included species of *Alternaria*, *Cladosporium*, *Epicoccum*, and *Sarocladium*. These genera include well-know saprobes and pathogens of taxonomically diverse plant species. Therefore, finding them as part of the mycobiota colonizing hardwood trees was not surprising.

Unlike for example, scolytine beetles (Curculionidae: Scolytinae), very little is known regarding the relationships between cerambycid beetles and fungi. Adults of these insects never enter the wood, and any organisms such as fungi carried on their bodies would need to be associated with oviposition. While these insects are well-known to transmit nematodes such as the devastating pine pathogen *Bursaphelenchus xylophilus* (pine wood nematode) during oviposition (Wingfield 1987; Akbulut and Stamps 2012), a fixed association between them and fungi is not known. In this regard, Wingfield et al. (2016) suggested that mites carried on the bodies of Cerambycidae could be implicated in the transmission of fungi found in their galleries. The same situation could be true also for *S. carcharias* and *A. glabripennis*, but further investigations would be needed to confirm it.

Species diversity of wood colonised by the insects was higher than intact wood highlighting the importance of larval tunnel for fungal diversity. We detected a number of pathogenic fungi capable of killing living cells, although many of these are considered as weak and opportunistic pathogens or endophytes. We also found several species with known potential of enzymatic degradation of wood components. The availability of such fungi closely associated with the insect suggests that fungal enzymes ingested by the larvae play an in key role in improved nutritional quality of the longhorn beetles. The study shows that cerambycid beetles are associated with a diverse microbial assemblage, which can participate in wood degradation by larvae either killing or breaking down host tissue.

Acknowledgements This study was financially supported by the University of Helsinki (RL); the Academy of Finland (FOA); the members of the Tree Protection Co-operative Programme (TPCP) and the THRIP initiative of the Department of Trade and Industry (RL, MJW), South Africa. We acknowledge the staff at the Finnish Food Safety Authority (Evira) for their support to this study.

Funding This study was funded by the University of Helsinki (RL); the Academy of Finland (FOA), the members of the Tree Protection Co-operative Programme (TPCP) and the THRIP initiative of the Department of Trade and Industry (RL, MJW), South Africa.

Compliance with ethical standards

Conflict of interest The authors declare that have no conflict of interest.

References

- Akbulut S, Stamps WT (2012) Insect vectors of the pinewood nematode: a review of the biology and ecology of *Monochamus* species. For Pathol 42:89–99. <https://doi.org/10.1111/j.1439-0329.2011.00733.x>
- Albrectsen BR, Björkén L, Varad A, Hagner Å, Wedin M, Karlsson J, Jansson S (2010) Endophytic fungi in European aspen (*Populus tremula*) leaves—diversity, detection, and a suggested correlation with herbivory resistance. Fungal Divers 41:17–28
- Ayayee P, Rosa C, Ferry JG, Felton GW, Saunders M, Hoover K (2014) Gut microbes contribute to nitrogen provisioning in a wood-feeding cerambycid. Environ Entomol 43:903–912. <https://doi.org/10.1603/EN14045>
- Blanchette RA, Held BW, Jurgens JA, McNew DL, Harrington TC, Duncan SM, Farrell RL (2004) Wood-destroying soft rot fungi in the historic expedition huts of Antarctica. Appl Environ Microbiol 3:1328–1335
- Casieri L, Hofstetter V, Viret O, Gindro K (2009) Fungal communities living in the wood of different cultivars of young *Vitis vinifera* plants. Phytopathol Mediterr 48:73–83. https://doi.org/10.14601/Phytopathol_Mediterr-2876
- Chararas C, Pignal M-C (1981) Étude du rôle de deux levures isolées dans le tube digestif de *Phoracantha semipunctata*, Coléoptère Cerambycidae xylophage spécifique des Eucalyptus. Comptes Rendus de l'Académie des Sciences de Paris 292:109–112
- Crous PW, Groenewald JZ, Gams W (2003) Eyspot of cereals revisited: ITS phylogeny reveals new species relationships. Eur J Plant Pathol 109:841–850. <https://doi.org/10.1023/A:1026111030426>
- Day MJ, Hall JC, Currah RS (2012) Phialide arrangement and character evolution in the helotiale anamorph genera *Cadophora* and *Phialocephala*. Mycologia 104:371–381. <https://doi.org/10.3852/11-059>
- Drenkhan T, Sibul I, Kasanen R, Vainio EJ (2013) Viruses of *Heterobasidion parviporum* persist within their fungal host during passage through the alimentary tract of *Hylobius abietis*. For Pathol 43:317–323
- Drenkhan T, Kasanen R, Vainio EJ (2016) *Phlebiopsis gigantea* and associated viruses survive passing through the digestive tract of *Hylobius abietis*. Biocontrol Sci Technol 26:320–330
- EPP0 (2016) Data sheets on quarantine pests—*Anoplophora glabripennis*. http://www.eppo.int/QUARANTINE/data_sheets/insects/ANOLGL_ds.pdf
- Francke-Grosman H (1967) Ectosymbiosis in wood-inhabiting insects. Symbiosis 2:141–205
- Gams W (2000) *Phialophora* and some similar morphologically little-differentiated anamorphs of divergent ascomycetes. Stud Mycol 45:187–199
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes—application to the identification of mycorrhiza and rusts. Mol Ecol 2:113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Geib SM, Scully ED, del Mar Jimene-Gasco M, Carlson JE, Tien M, Hoover K (2012) Phylogenetic analysis of *Fusarium solani* associated with the Asian longhorned

- beetle, *Anoplophora glabripennis*. Insects 3:141–160. <https://doi.org/10.3390/insects3010141>
- Gibson CM, Hunter MS (2009) Extraordinary widespread and fantastically complex: comparative biology of endosymbiotic bacterial and fungal mutualists of insects. Ecol Lett 13:223–234
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. Appl Environ Microbiol 61:1323–1330
- Gramaje D, Mostert L, Armengol J (2011) Characterization of *Cadophora luteo-olivacea* and *C. melinii* isolates obtained from grapevines and environmental samples from grapevine nurseries in Spain. Phytopathol Mediterr 50:112–126. https://doi.org/10.14601/Phytopathol_Mediterr-8723
- Greenleaf MA, Korf RP (1980) *Mollisia* in Macaronesia: an exercise in frustration. Mycotaxon 10:459–472
- Grünwald S, Pilhofer M, Höll W (2010) Microbial associations in gut systems of wood- and bark-inhabiting longhorned beetles [Coleoptera: Cerambycidae]. Syst Appl Microbiol 33:25–34
- Guindon S, Dufayard JF, Lefort V, Anisoma H, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59:207–321. <https://doi.org/10.1093/sysbio/syq010>
- Haack RA, Cavey JF, Hoebeke ER, Law K (1996) *Anoplophora glabripennis*: a new tree-infesting exotic cerambycid invades New York. News Mich Entomol Soc 41:1–3
- Haack RA, Hérard F, Sun J, Turgeon JJ (2010) Managing invasive populations of Asian longhorned beetle and citrus longhorned beetle: a worldwide perspective. Annu Rev Entomol 55:521–546. <https://doi.org/10.1146/annurev-ento-112408-085427>
- Hallaksela A-M (1999) Lahoisuus haavikossa. Metsäntutkimuslaitoksen tiedonantoja 725
- Halleen F, Mostert L, Crous PW (2007) Pathogenicity testing of lesser-known vascular fungi of grapevines. Australas Plant Pathol 36:277–285. <https://doi.org/10.1071/AP07019>
- Harrington TC, McNew DL (2003) Phylogenetic analysis places the *Phialophora*-like anamorph genus *Cadophora* in the Helotiales. Mycotaxon 87:141–151
- Heliövaara K, Mannerkoski I, Siitonen J (2004) Suomen sarvijäärät. Longhorn beetles in Finland (Coleoptera, Cerambycidae). Tremex Press, Helsinki
- Hérard F, Ciampitti M, Maspero M, Krehan H, Benker U, Boegel C, Schrage R, Bouhot-Delduc L, Bialooki P (2006) *Anoplophora* species in Europe: infestations and management processes. EPPO Bull 36:470–474. <https://doi.org/10.1111/j.1365-2338.2006.01046.x>
- Hérard F, Maspero M, Ramualde N, Jucker C, Colombo M, Ciampitti M, Cavagna B (2009) *Anoplophora glabripennis* infestation (Col.: Cerambycidae) in Italy. EPPO Bull 39: 146–152. <https://doi.org/10.1111/j.1365-2338.2009.02286.x>
- Herr JR, Scully ED, Geib SM, Hoover K, Carlson JE, Geiser DM (2016) Genome sequence of *Fusarium* isolate MYA-4552 from the midgut of *Anoplophora glabripennis*, an invasive, wood-boring beetle. Genome Announc 4:e00544-16. <https://doi.org/10.1128/genomeA.00544-16>
- Hu J, Angeli S, Schuetz S, Luo Y, Hajek AE (2009) Ecology and management of exotic and endemic Asian longhorned beetle *Anoplophora glabripennis*. Agric For Entomol 11:359–375. <https://doi.org/10.1111/j.1461-9563.2009.00443.x>
- Hulcr J, Dunn RR (2011) The sudden emergence of pathogenicity in insect–fungus symbioses threatens naive forest ecosystems. Proc R Soc B 278:2866–2873. <https://doi.org/10.1098/rspb.2011.1130>
- Humble LM, Allen EA (2006) Forest biosecurity: alien invasive species and vectored organisms. Can J Plant Pathol 28:S256–S269. <https://doi.org/10.1080/0706060609507383>
- Jones KG, Dowd PF, Blackwell M (1999) Polyphyletic origins of yeast-like endocytobionts from anobiid and cerambycid beetles. Mycol Res 103:542–546. <https://doi.org/10.1017/S0953756298007308>
- Jurzitza G (1959) Physiologische Untersuchungen an Cerambycidensymbionten. Arch Mikrobiol 33:305–332
- Kasson MT, O'Donnell K, Rooney AP, Sink S, Ploetz JN et al (2013) An inordinate fondness for *Fusarium*: phylogenetic diversity of fusaria cultivated by ambrosia beetles in the genus *Euwallacea* on avocado and other plant hosts. Fungal Genet Biol 56:147–157. <https://doi.org/10.1016/j.fgb.2013.04.004>
- Katoh K, Standley D (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evol 30:772–780. <https://doi.org/10.1093/molbev/mst010.pmid:23329690>
- Kukor JJ, Martin MM (1986) Cellulose digestion in *Monochamus marmorator* Kby. (Coleoptera: Cerambycidae): role of acquired fungal enzymes. J Chem Ecol 12:1057–1070. <https://doi.org/10.1007/BF01638996>
- Kukor JJ, Cowan DD, Martin MM (1988) The role of ingested fungal enzymes in cellulose digestion in the larvae of Cerambycid beetles. Physiol Biochem Zool 61:364–371
- Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. Mol Biol Evol 33:1870–1877. <https://doi.org/10.1093/molbev/msw054>
- Lagerberg T, Lundberg G, Melin E (1927) Biological and practical researches into blueing in pine and spruce. Sven Skogsvardsforen Tidskr 25:145–272
- Lefort V, Longueville J-E, Gascuel O (2017) SMS: smart model selection in PhyML. Mol Biol Evol 34:2422–2424. <https://doi.org/10.1093/molbev/msx149>
- Lingafelter SW, Hoebeke ER (2002) Revision of *Anoplophora* (Coleoptera: Cerambycidae). Entomological Society of Washington, Washington
- Linnakoski R, Jankowiak R, Villari C, Kirisits T, Solheim H, de Beer ZW, Wingfield MJ (2016) The *Ophiostoma clavatum* species complex: a newly defined group in the Ophiostomatales including three novel taxa. Antonie Van Lee J Microb 109:987–1018. <https://doi.org/10.1007/s10482-016-0700-y>
- Lu M, Wingfield MJ, Gillette NE, Sun J-H (2011) Do novel genotypes drive the success of an invasive bark beetle–fungus complex? Implications for potential reinvasion. Ecology 92:2013–2019. <https://doi.org/10.1890/11-0687.1>
- Morales-Ramos JA, Rojas MG, Sittertz-Bhatkar H, Saldana G (2000) Symbiotic relationship between *Hypothenemus hampei* (Coleoptera: Scolytidae) and *Fusarium solani* (Moniliales: Tuberculariaceae). Ann Entomol Soc Am 93:541–547. [https://doi.org/10.1603/0013-8746\(2000\)093\[0541:SRBHHC\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2000)093[0541:SRBHHC]2.0.CO;2)



- Nardon P, Grenier AM (1989) Endocytobiosis in Coleoptera: biological, biochemical, and genetic aspects. In: Schwenmiller W, Gassner G (eds) Insect endocytobiosis. CRC Press, Boca Raton, pp 175–215
- Navarrete F, Abreo E, Martinez S, Bettucci L, Sandra L (2011) Pathogenicity and molecular detection of Uruguayan isolates of *Greeneria uvicola* and *Cadophora luteo-olivacea* associated with grapevine trunk diseases. *Phytopathol Mediterr* 50:166–175. https://doi.org/10.14601/Phytopathol_Mediterr-9188
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University, Uppsala
- O'Donnel K, Cigelnik E (1997) Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Mol Phylogenet Evol* 7:103–116. <https://doi.org/10.1006/mpev.1996.0376>
- Pärtel K (2016) Application of untrastructural and molecular data in the helotialen fungi. *Dissertationes Biologicae Universitatis Tartuensia* 304
- Ploetz RC, Hulcr J, Wingfield MJ, de Beer ZW (2013) Destructive tree diseases associated with ambrosia and bark beetles: black swan events in tree pathology? *Plant Dis* 97:856–872. <https://doi.org/10.1094/PDIS-01-13-0056-FE>
- Ran B, Huang S, Henriksson G (2016) Isolation of exceedingly low oxygen consuming fungal strains able to utilize lignin as carbon source. *Cellul Chem Technol* 50:811–817
- Riba G (1977) Étude ultrastructurale de la multiplication et de la dégénérescence des symbionts des larves de *Criocephalus rusticus* [Coleoptera: Cerambycidae]; influence du jeune. *Annales de la Société Entomologique de France (Nouvelle Série)* 13:153–157
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Sánchez Márquez S, Bills GF, Zabalgoceazcoa I (2007) The endophytic mycobiota of the grass *Dactylis glomerata*. *Fungal Divers* 27:171–195
- Santamaría O, Diez JJ (2005) Fungi in leaves, twigs and stem bark of *Populus tremula* from northern Spain. *For Pathol* 35:95–104
- Santini A, Faccoli M (2014) Dutch elm disease and elm bark beetles: a century of association. *iForest* 8:126–134. <https://doi.org/10.3832/ifor1231-008>
- Schloss PD, Delalibera I Jr, Handelsman J, Raffa KF (2006) Bacteria associated with the guts of two wood-boring beetles: *Anoplophora glabripennis* and *Saperda vestita* (Coleoptera). *Environ Entomol* 35:625–629
- Scully ED, Geib SM, Hoover K, Tien M, Tringe SG, Barry KW, Glavina del Rio T, Chovatia M, Herr JR, Carlson JE (2013) Metagenomic profiling reveals lignocellulose degrading system in a microbial community associated with a wood-feeding beetle. *PLoS ONE* 8:e73827. <https://doi.org/10.1371/journal.pone.0073827>
- Scully ED, Geib SM, Carlson JE, Tien M, McKenna D, Hoover K (2014) Functional genomics and microbiome profiling of the Asian longhorned beetle (*Anoplophora glabripennis*) reveal insights into the digestive physiology and nutritional ecology of wood feeding beetles. *BMC Genom* 15:1096. <https://doi.org/10.1186/1471-2164-15-1096>
- Stenlid J, Oliva J (2016) Phenotypic interactions between tree hosts and invasive forest pathogens in the light of globalization and climate change. *Philos Trans R Soc B* 371:20150455. <https://doi.org/10.1098/rstb.2015.0455>
- Sugiura N (1978) Further analysis of the data by Akaike's information criterion and the finite corrections. *Commun Stat Theory Methods* 7:13–26
- Swofford DL (2002) PAUP* 4.0: phylogenetic analysis using parsimony (* and other methods). Sinauer Associates, Sunderland
- Terhonen E, Marco T, Sun H, Jalkanen R, Kasanen R, Vuorinen M, Asiegbu F (2011) The effect of latitude, season and needle-age on the mycobiota of scots pine (*Pinus sylvestris*) in Finland. *Silva Fenn* 45:301–317. <https://doi.org/10.14214/sf.1320>
- Tomiczek C, Krehan H, Menschhorn P (2002) Dangerous Asian Longhorn Beetle found in Austria: a new threat for our trees? *AFZ/Der Wald Allgemeine Forst Zeitschrift für Waldwirtschaft und Umweltversorge* 57:52–54 (in German)
- Travadon R, Lawrence DP, Rooney-Latham S, Gubler WD, Wilcox WF, Rolshausen PE, Baumgartner K (2015) *Cadophora* species associated with wood-decay of grapevine in North America. *Fungal Biol* 119:53–66. <https://doi.org/10.1016/j.funbio.2014.11.002>
- Úrbez-Torres JR, Haag P, Bowen P, O'Gorman DT (2014) Grapevine trunk diseases in British Columbia: incidence and characterization of the fungal pathogens associated with Esca and Petri Diseases of grapevine. *Plant Dis* 98:469–482. <https://doi.org/10.1094/PDIS-05-13-0523-RE>
- Uvarov B (1929) Insect nutrition and metabolism. *Trans R Entomol Soc Lond* 76:255–343
- Välimäki S, Heliövaara K (2007) Hybrid aspen is not preferred by the large poplar borer (*Saperda carcharias*). *Arthropod Plant Interact* 1:205–211
- Vega FE, Blackwell M (2005) Insect-fungal associations: ecology and evolution. Oxford University Press, New York
- Watanabe H, Tokuda G (2010) Cellulosic systems in insects. *Annu Rev Entomol* 55:609–632. <https://doi.org/10.1146/annurev-ento-112408-085319>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR protocols: a guide to methods and applications. Academic Press, Millbrae, pp 315–321
- Wingfield MJ (1987) Fungi associated with the pine wood nematode, *Bursaphelenchus xylophilus*, and cerambycid beetles in Wisconsin. *Mycologia* 79:325–328
- Wingfield MJ, Garnas JR, Hajek A, Hurley BP, de Beer ZW, Taerum SJ (2016) Novel and co-evolved associations between insects and microorganisms as drivers of forest pestilence. *Biol Invasions* 18:1045–1056. <https://doi.org/10.1007/s10530-016-1084-7>



Journal : **10482**

Article : **1112**

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details Required	Author's Response
AQ1	Please check and confirm the corresponding affiliation is correctly identified.	
AQ2	Katoh et al. (2013) has been changed to Katoh and Standley (2013) so that this citation matches the list.	
AQ3	Akbulut et al. (2012) has been changed to Akbulut and Stamps (2012) so that this citation matches the list.	
AQ4	Wingfield et al. (1987) has been changed to Wingfield (1987) so that this citation matches the list.	
AQ5	Please provide Accessed date for the reference EPPO (2016).	
AQ6	Please provide page ranges for the reference Hallaksela (1999).	